

Living With Sarcoidosis


Sarcoidosis is a disease associated with **inflammation**, a response of cells to injury. The cause of injury to cells affected by sarcoidosis is unknown. It can affect almost any organ, but usually starts in the lungs or lymph nodes. Sarcoidosis can occur in anyone at any age, but is more common among individuals aged 20 to 40 years. It affects black individuals more commonly than white. Sarcoidosis is not contagious and is rarely fatal.

The March 13, 2002, issue of JAMA includes an article about treating sarcoidosis affecting the lungs with **corticosteroids**, a type of medication that reduces inflammation.

SYMPTOMS

Sarcoidosis usually develops slowly, although symptoms can appear suddenly and then disappear. The disease may be diagnosed by evaluating the patient's medical history, physical examination, and chest x-ray and eliminating other possible causes for the symptoms. Sarcoidosis may affect a variety of different organs. The lungs and lymph nodes are most commonly involved. Symptoms range from mild to severe. General symptoms may include fatigue, fever, and weight loss.

Symptoms in Patients With Sarcoidosis



FOR MORE INFORMATION

- National Heart, Lung, and Blood Institute
301/592-8573
www.nhlbi.nih.gov
- American Lung Association
212/315-8700
www.lungusa.org

INFORM YOURSELF

To find this and previous JAMA Patient Pages, go to the Patient Page Index on JAMA's Web site at www.jama.com.

Sources: American Academy of Family Physicians, Arthritis Foundation, Mayo Foundation, National Heart, Lung, and Blood Institute, National Sarcoidosis Resource Center, Vanderbilt Medical Center

TREATMENT

Sarcoidosis may go away without any treatment, and about 50% of people who get it recover completely. Depending on the severity of your symptoms, your doctor may either monitor you for several months or prescribe medication to treat the symptoms. The most common medications for sarcoidosis are corticosteroids, such as prednisone. Corticosteroids are often helpful in reducing symptoms, but may cause side effects such as mood swings, weight gain, increased blood pressure, and stomach ulcers. Therefore, their use should be carefully monitored.

Although much about sarcoidosis remains unknown, it is usually not disabling and often goes away in 2 to 3 years. People who have the disease for longer can usually lead normal lives.

Lise M. Stevens, MA, Writer

Cassio Lynn, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be reproduced noncommercially by physicians and other health care professionals to share with patients. Any other reproduction is subject to AMA approval. To purchase bulk reprints, call 718/946-7424.

